

PAPER NAME AUTHOR

The Relationship Of Characteristics Of M aternal With The Duration Of Placenta Re moval In The Bio Cli

WORD COUNT CHARACTER COUNT

2449 Words 12511 Characters

PAGE COUNT FILE SIZE

5 Pages 270.7KB

SUBMISSION DATE REPORT DATE

Apr 18, 2023 7:47 PM GMT+7 Apr 18, 2023 7:47 PM GMT+7

Yanti

# 23% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

• 15% Publications database

Crossref database

• Crossref Posted Content database

• 17% Submitted Works database

# Excluded from Similarity Report

Internet database

• Bibliographic material



#### Journal homepage: https://pcijournal.org/index.php/iamsph

## **International Archives of Medical Sciences and Public Health**



Published: Pena Cendekia Insani

# The Relationship Of Characteristics Of Maternal With The Duration Of Placenta Removal In The Bio Clinic In 2022

Yanti<sup>1</sup>, Afrida Yelni<sup>2</sup>, Faradita Wahyuni<sup>3</sup>

<sup>1,2</sup> Departmen of Wifery, STIKES Senior Medan

#### Article Info

#### Article history:

Received: 28 March 2022 Revised: 27 April 2022 Accepted: 30 June 2022

#### Keywords:

Age, Parity and Placenta Release

#### **ABSTRACT**

Hemorrhage with retained placenta accounts for a small proportion of cases of postpartum hemorrhage, but the incidence of retained placenta accounts for the maternal mortality rate due to retained placenta and the incidence is 0.8-1.2% for each birth. The cause of retained placenta between the placenta has not been separated from the uterine wall and the placenta has separated, but has not been born. Sased on the above background, the formulation of the problem in this study is how the characteristics of the mother in labor and the duration of placental separation at the Biological Clinic in 2022. The research design used in this study was an analytical survey with a cross sectional approach. This done to explore how and why the phenomenon occurs, and to study he relationship between the independent variable and the dependent variable. Population is the whole object of research or object under study. In this study, the population was all mothers who gave birth at the Biological Clinic as many as 153 people. Sampling in this study using the Slovin formula as many as 60 people. With Random Sampling Technique. The results of statistical tests with 130 square at = 0.05 Technique. The results of statistical tests with  $\frac{1}{12}$  is square at = 0.05 obtained P value = 0.006 (p> $\alpha$ ), thus showing that there is a significant relationship between maternal age and the duration of placental separation. The results of statistical tests with square at = 0.05 separation. The results of statistical tests with his-square at = 0.05 obtained P value = 0.023 (p> $\alpha$ ), thus showing that there is a significant relationship between maternal parity and the duration of placental separation. Conclusion In this study, it can be seen that there is a relationship between maternal characteristics and the duration of placental separation at the Biological Clinic in 2022. It is recommended as input so that future researchers can make this research a comparison with previous research and it is hoped that researchers will be able to develop this study with variables that different.

<sup>5</sup>his is an open access article under the <u>CC BY-SA</u>license.



#### Corresponding Author:

Yanti.

Department Of Wifery STIKES Senior Medan

Email: afriwanyanti@gmail.com

#### 1. INTRODUCTION

Pregnancy and childbirth have great health risks, including for women who have no previous health problems. Postpartum hemorrhage is bleeding of more than 500-600 ml within 24 hours after the child is born. Usually after the fetus is born, a few minutes later the process of releasing the placenta begins with a little bleeding.

Based on WHO (World Health Organization) data worldwide, there are 500,000 maternal deaths per year to pregnancy-related causes, 99% of these deaths occur in developing countries. In developing countries in general, the maternal mortality ratio ranges from 100-1000 per 100,000 live births

Based on reports from district/city profiles, the Maternal Mortality Rate <sup>10</sup>IMR) reported in North Sumatra in 2020 is only 75/100,000 live births, but this cannot yet describe the actual MMR in the population. Based on the results of the 2020 population census, the <sup>4</sup>IMR in North Sumatra is 328/100,000 live births, this figure is still quite high when compared to the national figure for the 2020 population census of 259/100,000 live births. <sup>4</sup>Dassed on the results of the MMR and Infant Mortality Rate (IMR) <sup>4</sup>Durvey conducted by the North Sumatra Provincial Health Office, it was 268 per 100,000 live births. Based on estimates, this maternal mortality rate will not decrease until 2020.

assed on the above background, the formulation of the problem in this study is how the characteristics of the mother in labor are related to the length of time the placenta is released at the Medan Biological Clinic in 2022.

## 2. RESEARCH METHODE

This is done to explore how and why the phenomenon occurs, and to study are relationship between the independent variable and the dependent variable. In this study age, parity, and early initiation of breastfeeding as independent variables and duration of placental separation as dependent variables. This research was conducted in a biological clinic in 2022. This research was conducted from January 2022 to July 2022. The population is the entire object of research or the object under study. In this study, the population is all maternity mothers who give birth at the Biological Clinic in 2022. A total of 153. The sample is part of the number and characteristics possessed by the population. While sampling means taking a sample or taking a part of the population or universe as a representative (representation). Sampling in this study used the slovin formula. So the sample obtained was 60 people, sampling using the Random Sampling technique

# 3. RESULT AND ANALYSIS

# **RESULT**

Table 1. Cross-tabulation of Maternal Age with Length of Placenta Release at the Biological Clinic in 2022

		Lan	ıa Pelepa	asan Pla	senta	т.	otal	_
No	Usia	≤ 15 M	<b>Ienit</b>	> 15 N	<b>Ienit</b>	10	iai	Asym.Sig (p)
		f	%	f	%	F	%	_
1	20-35 Tahun	22	36,7	14	23,3	39	60	0.006
2	<20 dan >35 Tahun	6	10	18	30	21	40	0,006
	Jumlah	28	46,7	32	53,3	60	100	

years (60%) with placental separation duration 15 minutes as many as 22 people (36.7%), and with placental separation duration > 15 minutes as many as 14 people (23.3%), while There were 24 people aged <20 and > 35 years (40%) with a placental separation duration of 15 minutes as many as 6 people (10%) and 15 people (30%). 3 ased on the test results using the Chi-Square analysis tool, it can be seen that the value = 0.006 <0.05 so that there is a relationship between maternal age and the duration of placental separation.

Table 2. Cross-tabulation of maternal parity with the duration of placental detachment at the Biological Clinic in 2022

		Lam	a Pelepa	asan P	lasenta	Т.	-4-1	A G!
No	<b>Paritas</b>	≤ 15	Menit	> 15	Menit	10	otal	Asym.Sig
		f	%	f	%	F	%	<b>-</b> ( <b>p</b> )
1	Primipara	16	26,7	9	15,0	25	41,7	0,023
2	Multipara	12	20,0	23	38,3	35	58,3	0,023
	Jumlah	28	46,7	32	53,3	60	100	

Tasked on Table 2, it can be seen that from 60 mothers (100%). There were 35 multiparous mothers (58.3%) with placental separation duration > 15 minutes as many as 23 people (38.3%), and with placental separation duration 15 minutes as many as 12 people (20%), while the primiparous mothers were 25 people. (41.7%) with placental separation duration 15 minutes as many as 16 people (26.7%) and placenta detachment duration > 15 minutes as many as 9 people (15%). Based on the statistical test using Chi-Square, it was found that p = 0.023 <0.05 so that there was a parity relationship with the duration of placental separation.

Table 3. Cross-tabulation of Pregnancy Distance with Length of Placenta Release at the Biological Clinic in 2022

No	Paritas			enta		Т	otal	Asym.Sig (p)
		<u>≤15</u>	Menit	> 15	Menit			_ (P)
10		f	<b>%</b>	f	<b>%</b>	$\mathbf{F}$	%	
	< 2 tahun	16	26,7	26	43,3	42	70	_
2	2-5 Tahun	12	20	5	8,3	17	28,3	0,049
3	> 5 Tahun	0	0	1	3,1	1	1,7	
	Jumlah	28	46,7	32	53,3	60	100	

 $^{20}$ ased on Table 3, it can be seen that from 60 mothers (100%). The mothers who had the most gestational interval < 2 years were 42 people (70%) with a duration of placental separation > 15 minutes as many as 26 people (43.3%), and with a duration of placental separation 15 minutes as many as 16 people (26.7%).), while those who had the least gestational interval > 5 years were 1 person (1.7%) with a length of placental separation > 15 minutes.

# **DISCUSSION**

The relationship between parity and the duration of placental detachment shows a significant relationship, this is evident from the results of studies that show the duration of placental detachment assed on parity. The results of statistical tests with Chi-square at = 0.05 obtained P

value = 0.023 (p> $\alpha$ ), thus showing that there is a significant relationship between maternal parity and the duration of placental separation.

This is in accordance with the theory which states that parity that is too high has a risk of childbirth, one of which is the detachment of the placenta. Multiparas have a longer time to deliver the placenta because in multiparas the uterus tends to work less efficiently at all stages of labor.

This is in line with Gita Kostania's research on the relationship between parity and the duration of placental release in maternity at BPS Sarwo Endah Kadipaten, Andong, Boyolali from January to April 2011 which add that there was a significant relationship between parity and the duration of placental detachment in maternity. Square p = 0.023.

Based on the results of the study, the authors concluded that there was a relationship between parity and the duration of placental separation, this is because parity greatly affects placental separation, especially in multiparas who have a high risk, because more and more mothers give birth, the condition of the uterus becomes less efficient, namely there is no attachment site for the placenta, because the former implantation in a previous delivery.

The relationship between the distance of pregnancy and the duration of placental separation shows a significant relationship, this is evident from the results of studies that show the length of time for placental separation based on the distance of pregnancy. The results of statistical tests with chi-square at = 0.05 obtained P value = 0.049 (p> $\alpha$ ), thus showing that there is a significant relationship between the distance between the mother's pregnancy and the duration of placental separation.

In accordance with the theory that a short pregnancy interval will directly have an effect on the health of women and the health of the fetus they contain. A woman after giving birth takes 2 to 3 years to recover her body and prepare herself for the next pregnancy and childbirth. The closer the distance between the mother's pregnancy, the more it will affect the incidence of retained placenta.

Based on the results of the study, the authors concluded that there was a relationship between the distance of pregnancy and the duration of placental separation, this is because the distance of pregnancy is too close resulting in high-risk labor, the more often pregnant women and childbirth, the closer the distance between pregnancy and delivery, the elasticity of the uterus is increasingly disturbed, as a result the uterus does not become pregnant. fully contracted.

# 4. CONCLUSION

2 ased on the results of research conducted with the title Relationship of Maternal Characteristics in Maternity with the Length of Placenta Release at the Hj Clinic. Dermawati Nasution Medan in 2016, it can be concluded as follows:

- 1) Of the 60 people studied, most of the mothers who gave birth aged  $\leq$  20 and  $\geq$  35 years were 18 (30%), mothers with multipara were 35 people (58.3%), and most of the mothers gave birth with a gestation interval of  $\leq$  2 years. namely 26 people (43.3%).
- 2) According to the category of duration of placental detachment at the Hj. Dermawati Nasution Medan In 2016, from 60 cases found 32 cases (53.3%) > 15 minutes and 28 cases (46.7%) < 15 minutes.
- 3) In this study, it can be seen that there is a relationship between maternal characteristics and the duration of placental separation at the Hj Clinic. Dermawati Nasution Medan 2016.

## References

- 1. Mochtar R. Sinopsis Obstetri. 3rd ed. Indra L, editor. Jakarta: EGC; 2012.
- Manuaba IAC, dkk. Ilmu Kebidanan, Penyakit Kandungan, dan KB. 2nd ed. Jakarta: Buku Kedokteran EGC; 2012.
- 3. RN, dkk. Usia Dan Paritas Dengan Plasenta Previa Pada Ibu Bersalin. 2012 juni.
- 4. DepKes RI. Profil Kesehatan Indonesia. [Online].; 2015 [cited 2017 January 26. Available from: <a href="http://www.search-document.com/pdf/sumut-profil-2014.html">http://www.search-document.com/pdf/sumut-profil-2014.html</a>.
- 5. DinKes Sumut. Profil Kesehatan Indonesia. [Online].; 2014 [cited 2016 November 26. Available from: <a href="http://www.search-document.com/pdf/sumut-profil-2014.html">http://www.search-document.com/pdf/sumut-profil-2014.html</a>.
- 6. Prawirohardjo S. Ilmu Kebidanan. 4th ed. Jakarta: Bina Pustaka Sarwono Prawirohardjo; 2014.
- 7. Agustin, Ika Minda; Atik Setyaningsih. Hubungan Umur Dan Paritas dengan Lamanya Pelepasan Plasenta. journal.stikeseub.ac.id. 2011 Juni; III(1).
- 8. H, M. Ilmu kebidanan patologi dan fisiologi persalinan Medika YE, editor. Yogyakarta; 2012.
- 9. Zeu E. Hubungan antara Umur dan Paritas Ibu dengan Kejadian Retensio Plasenta di Puskesmas Jagir Surabaya Tahun 2012. 2012.
- 10. Maryunani A. Kegawatdaruratan Maternal dan Neonatal Terpadu Jakarta: Trans Info Media; 2015.
- 11. Intan M. Hubungan Paritas dengan Retensio Plasenta pada Ibu Bersalin di RSUD Syekh Yusuf Kabupaten Gowa. 2012.
- 12. Kostania G, Purnamasari DE. Hubungan Paritas dengan Lamanya Pelepasan Plasenta pada Ibu Bersalin di BPS Sarwo Endah Kadipaten, Andong, Boyolali. 2011 Januari-April.
- 13. E. Asuhan Kebidanan pada Persalinan Patologi Yogyakarta: Pustaka Pelajar; 2013.
- 14. Yeyeh A. Asuhan Kebidanan IV (patologi) Jakarta: Trans Info Media; 2015.
- 15. Mochtar R. Sinopsis Obstetri. 3rd ed. Jakarta: Buku Kedokteran EGC; 2013.
- 16. Rukiyah AY. Asuhan Kebidanan IV (Patologi). 2nd ed. Jakarta: CV. Trans Info Media; 2015.
- 17. Fisiologi Plasenta dalam Kehamilan. [Online].; 2014 [cited 2014 Desember. Available from: <a href="http://wawasankehamilan.blogspot.co.id/2014/12/fisiologi-plasenta-dalam-kehamilan.html?m=1">http://wawasankehamilan.blogspot.co.id/2014/12/fisiologi-plasenta-dalam-kehamilan.html?m=1</a>.
- 18. Wahyuni A. Kala III Pengawasan. [Online].; 2016 [cited 2016 februari 16. Available from: <a href="http://www.academia.edu/9329540/Kala\_III\_Pengawasan">http://www.academia.edu/9329540/Kala\_III\_Pengawasan</a>.
- 19. Manajemen Aktif Kala III. [Online]. [cited 2011 Oktober 25. Available from: <a href="http://www.lusa.web.id/manajemen-aktif-kala-iii/">http://www.lusa.web.id/manajemen-aktif-kala-iii/</a>.
- 20. Wawan A MD. Teori & Pengukuran Pengetahuan, Sikap dan Perilaku Manusia. 2nd ed. Yogyakarta: Nuha Medika; 2016.
- 21. Rismalinda. Buku Ajar Asuhan Kebidanan Kehamilan. 1st ed. T I, editor. Jakarta Timur: CV. Trans Info Media; 2015.
- 22. Maryunani A. Buku Praktis Kehamilan dan Persalinan Patologis (Risiko Tinggi dan Komplikasi) dalam kebidanan Jakarta: Trans Info Media; 2016.
- 23. Muhammad I. Panduan Penyusunan Karya Tulis Ilmiah Bidang Kesehatan Menggunakan Metode Ilmiah. 5th ed. RB S, editor. Medan: Citapustaka Media Perintis; 2015.



# 23% Overall Similarity

Top sources found in the following databases:

- 15% Publications database
- Crossref Posted Content database
- · Crossref database
- 17% Submitted Works database

# **TOP SOURCES**

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

Universitas Sebelas Maret on 2023-03-24 Submitted works	2%
Pertiwi Perwiraningtyas, Anita Rahmawati. "Factor Analysis of Caesare  Crossref	2%
Muhammad Nur Abdullah, Arifin Ahmad, Hamsu Abdul Gani. "Develop  Crossref	2%
Diah Kusuma Dewi, Purhadi, Sutikno. "Geographically Weighted Bivari  Crossref	. 2%
UC, Irvine on 2021-11-04 Submitted works	1%
Brown University on 2021-11-14 Submitted works	1%
Endriyani Syafitri, Suyanti Suwardi. "Faktor-Faktor yang Berhubungan d Crossref	1%
Dwiana Kartika Putri. "Hubungan Pengetahuan dan Sikap Ibu Hamil Tri  Crossref	1%
Universiti Teknologi MARA on 2021-08-19 Submitted works	1%



Badan PPSDM Kesehatan Kementerian Kesehatan on 2 Submitted works	2022-10-25
Sri Juliani. "Hubungan Dukungan Suami dan Motivasi B <sup>Crossref</sup>	idan dengan Ke
Universitas Nasional on 2021-01-12 Submitted works	•
Universitas Pamulang on 2023-03-03 Submitted works	•
Badan PPSDM Kesehatan Kementerian Kesehatan on 2 Submitted works	2022-09-15
Linda Zuliana, Haerawati Idris, Zulkarnain Zulkarnain. "I	Evaluasi Penyel
Meadowvale High School on 2002-03-27 Submitted works	•
S Jumiyati, N Possumah, A Hadid, S Bachri, P Dua. "Eco Crossref	o-friendly utilizat
Universitas Jenderal Soedirman on 2020-11-17 Submitted works	•
iGroup on 2014-10-20 Submitted works	
Siti Umi Kalsum, G M Saragih, Misbahul Munir. "Spatial Crossref	based mapping
Mansoura University on 2020-08-15 Submitted works	